



## **General Certificate of Education**

# **Mathematics 6360**

**MPC1      Pure Core 1**

## **Mark Scheme**

*2007 examination - June series*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: [www.aqa.org.uk](http://www.aqa.org.uk)

Copyright © 2007 AQA and its licensors. All rights reserved.

#### COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

---

### Key to mark scheme and abbreviations used in marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
√ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A <sub>2,1</sub>	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

### No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**


## MPC1

Q	Solution	Marks	Total	Comments
1(a)(i)	Gradient $AB = \frac{-1-5}{6-2}$ or $\frac{5-1}{2-6}$	M1		$\pm \frac{6}{4}$ implies M1
	$= \frac{-6}{4} = -\frac{3}{2}$	A1	2	AG
(ii)	$\left. \begin{matrix} y-5 \\ y+1 \end{matrix} \right\} = -\frac{3}{2} \left\{ \begin{matrix} (x-2) \\ (x-6) \end{matrix} \right.$	M1		or $y = -\frac{3}{2}x + c$ and attempt to find $c$
	$\Rightarrow 3x + 2y = 16$	A1	2	OE; must have integer coefficients
(b)(i)	Gradient of perpendicular = $\frac{2}{3}$	M1		or use of $m_1m_2 = -1$
	$\Rightarrow y - 5 = \frac{2}{3}(x - 2)$	A1	2	$3y - 2x = 11$ (no misreads permitted)
(ii)	Substitute $x = k, y = 7$ into their (b)(i)	M1		or grads $\frac{7-5}{k-2} \times \frac{-3}{2} = -1$
	$\Rightarrow 2 = \frac{2}{3}(k - 2) \Rightarrow k = 5$	A1	2	or Pythagoras $(k - 2)^2 = (k - 6)^2 + 8$
<b>Total</b>			<b>8</b>	
2(a)	$\frac{\sqrt{63}}{3} = \sqrt{7}$ or $\frac{3\sqrt{7}}{3}$	B1		or $\frac{(\sqrt{7}\sqrt{63} + 14 \times 3)}{3\sqrt{7}}$
	$\frac{14}{\sqrt{7}} = 2\sqrt{7}$ or $\frac{14\sqrt{7}}{7}$	B1		or $\frac{\sqrt{7}}{\sqrt{7}} ( \quad )$ M1
	$\Rightarrow \text{sum} = 3\sqrt{7}$	B1	3	$\Rightarrow$ correct answer with all working correct A2
(b)	Multiply by $\frac{\sqrt{7}+2}{\sqrt{7}+2}$	M1		
	Denominator = $7 - 4 = 3$	A1		
	Numerator = $(\sqrt{7})^2 + \sqrt{7} + 2\sqrt{7} + 2$	m1		multiplied out (allow one slip) $9 + 3\sqrt{7}$
	Answer = $\sqrt{7} + 3$	A1	4	
<b>Total</b>			<b>7</b>	


## MPC1 (cont)

Q	Solution	Marks	Total	Comments
<b>3(a)(i)</b>	$(x+5)^2$ -6	B1 B1	2	$p = 5$ $q = -6$
<b>(ii)</b>	$x_{\text{vertex}} = -5$ (or their $-p$ ) $y_{\text{vertex}} = -6$ (or their $q$ )	B1✓ B1✓	2	may differentiate but must have $x = -5$ and $y = -6$ . Vertex $(-5, -6)$
<b>(iii)</b>	$x = -5$	B1	1	
<b>(iv)</b>	Translation (not shift, move etc) through $\begin{bmatrix} -5 \\ -6 \end{bmatrix}$ (or 5 left, 6 down)	E1 M1 A1	3	and NO other transformation stated either component correct M1, A1 independent of E mark
<b>(b)</b>	$x+11 = x^2 + 10x + 19$ $\Rightarrow x^2 + 9x + 8 = 0$ or $y^2 - 13y + 30 = 0$ $(x+8)(x+1) = 0$ or $(y-3)(y-10) = 0$ $x = -1$ } or $x = -8$ } $y = 10$ } $y = 3$ }	M1 m1 A1 A1	4	quadratic with all terms on one side of equation attempt at formula (1 slip) or to factorise both $x$ values correct both $y$ values correct and linked SC $(-1, 10)$ B2, $(-8, 3)$ B2 no working
<b>Total</b>			<b>12</b>	
<b>4(a)(i)</b>	$t^3 - 52t + 96$	M1 A1 A1	3	one term correct another term correct all correct (no + c etc)
<b>(ii)</b>	$3t^2 - 52$	M1 A1✓	2	ft one term correct ft all "correct"
<b>(b)</b>	$\frac{dy}{dt} = 8 - 104 + 96$ $= 0 \Rightarrow$ stationary value Substitute $t = 2$ into $\frac{d^2y}{dt^2}$ ( $= -40$ ) $\frac{d^2y}{dt^2} < 0 \Rightarrow$ max value	M1 A1 M1 A1	4	substitute $t = 2$ into their $\frac{dy}{dt}$ CSO; shown = 0 + statement any appropriate test, e.g. $y'(1)$ and $y'(3)$ all values (if stated) must be correct
<b>(c)</b>	Substitute $t = 1$ into their $\frac{dy}{dt}$ Rate of change = $45 \text{ (cms}^{-1}\text{)}$	M1 A1✓	2	must be their $\frac{dy}{dt}$ NOT $\frac{d^2y}{dt^2}$ ft their $y'(1)$
<b>(d)</b>	Substitute $t = 3$ into their $\frac{dy}{dt}$ $(27 - 156 + 96 = -33 < 0)$ $\Rightarrow$ decreasing when $t = 3$	M1 E1✓	2	interpreting their value of $\frac{dy}{dt}$ allow increasing if their $\frac{dy}{dt} > 0$
<b>Total</b>			<b>13</b>	

## MPC1 (cont)

Q	Solution	Marks	Total	Comments
5(a)(i)	Centre $(-3, 2)$	M1	2	$\pm 3$ or $\pm 2$
		A1		correct
(ii)	Radius = 5	B1	1	accept $\sqrt{25}$ but not $\pm\sqrt{25}$
(b)(i)	$3^2 + (-4)^2 = 9 + 16 = 25$ $\Rightarrow N$ lies on circle	B1	1	must have $9 + 16 = 25$ or a statement
(ii)		M1	2	must draw axes; fit their centre in correct quadrant
		A1		correct (reasonable freehand circle enclosing origin)
(iii)	Attempt at gradient of $CN$	M1	3	withhold if subsequently finds tangent
	$\text{grad } CN = -\frac{4}{3}$	A1		CSO
	$y = -\frac{4}{3}x - 2$ (or equivalent)	A1✓		fit their grad $CN$
(c)(i)	$P(2, 6)$ Hence $PC^2 = 5^2 + 4^2$ $\Rightarrow PC = \sqrt{41}$	M1	2	“their” $PC^2$
		A1		
(ii)	Use of Pythagoras correctly $PT^2 = PC^2 - r^2 = 41 - 25$ , where $T$ is a point of contact of tangent $\Rightarrow PT = 4$	M1	3	fit their $PC^2$ and $r^2$
		A1✓		<b>Alternative</b> sketch with vertical tangent M1 showing that tangent touches circle at point $(2, 2)$ A1
		A1		hence $PT = 4$ A1
<b>Total</b>			<b>14</b>	

## MPC1 (cont)

Q	Solution	Marks	Total	Comments
6(a)(i)	$f(1) = 1 + 4 - 5$	M1	2	must find $f(1)$ NOT long division shown = 0 plus a statement
	$\Rightarrow f(1) = 0 \Rightarrow (x-1)$ is factor	A1		
(ii)	Attempt at $x^2 + x + 5$	M1	2	long division leading to $x^2 \pm x + \dots$ or equating coefficients $p = 1, q = 5$ by inspection scores B1, B1
	$f(x) = (x-1)(x^2 + x + 5)$	A1		
(iii)	$(x =) 1$ is real root	B1	3	not the cubic! CSO; all values correct plus a statement
	Consider $b^2 - 4ac$ for their $x^2 + x + 5$	M1		
	$b^2 - 4ac = 1^2 - 4 \times 5 = -19 < 0$ Hence no real roots (or only real root is 1)	A1		
(b)(i)	$\int \dots dx = \frac{x^4}{4} + 2x^2 - 5x (+c)$	M1 A1 A1	3	one term correct unsimplified second term correct unsimplified all correct unsimplified
	$[4 + 8 - 10] - \left[ \frac{1}{4} + 2 - 5 \right]$	M1		
	$= 4 \frac{3}{4}$ Area of $\Delta = \frac{1}{2} \times 11 = 5 \frac{1}{2}$ $\Rightarrow$ shaded area $= 5 \frac{1}{2} - 4 \frac{3}{4}$ $= \frac{3}{4}$	A1 B1 A1		
	<b>Total</b>		<b>14</b>	
7(a)	$b^2 - 4ac = 4 - 4(k-1)(2k-3)$	M1	3	(or seen in formula) condone one slip must involve $f(k) \geq 0$ (usually M1 must be earned)  at least one step of working justifying $\leq 0$ AG
	Real roots when $b^2 - 4ac \geq 0$	E1		
	$4 - 4(2k^2 - 5k + 3) \geq 0$ $\Rightarrow -2k^2 + 5k - 3 + 1 \geq 0$ $\Rightarrow 2k^2 - 5k + 2 \leq 0$	A1		
(b)(i)	$(2k-1)(k-2)$	B1	1	
(ii)	(Critical values) $\frac{1}{2}$ and 2	B1 $\checkmark$	3	fit their factors or correct values seen on diagram, sketch or inequality or stated  use of sketch / sign diagram M1A0 for $0.5 < k < 2$ or $k \geq 0.5, k \leq 2$
		M1		
	$\Rightarrow 0.5 \leq k \leq 2$	A1		
	<b>Total</b>		<b>7</b>	
	<b>TOTAL</b>		<b>75</b>	