



GCE AS/A level

0985/01

**MATHEMATICS S3
STATISTICS 3**

P.M. FRIDAY, 22 June 2012

1½ hours

ADDITIONAL MATERIALS

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator;
- statistical tables (Murdoch and Barnes or RND/WJEC Publications)

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen.

Answer **all** questions.

Sufficient working must be shown to demonstrate the **mathematical** method employed.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

1. Three numbers are chosen at random **without replacement** from the set $\{1, 2, 3, 4, 5\}$. Determine the sampling distribution of

(a) the mean of the three chosen numbers, [5]

(b) the median of the three chosen numbers. [2]

2. The manager of a factory that manufactures a certain type of string claims that its mean breaking strength is 100 Newtons. In order to test this claim, the breaking strengths, in Newtons, of a random sample of ten pieces of this string were determined with the following results.

97.1 100.3 98.6 97.7 101.2 97.6 98.9 101.1 98.5 99.3

You may assume that this is a random sample from a normal distribution with mean μ and variance σ^2 .

(a) Calculate unbiased estimates of μ and σ^2 . [4]

(b) (i) State suitable hypotheses for testing the manager's claim using a two-sided test.

(ii) Carry out an appropriate test at the 5% significance level. Giving a reason, state your conclusion in context. [7]

3. Alan plays a certain game on his computer and he wants to estimate the probability p of winning. He plays the game 120 times and wins 54 of these games. It may be assumed that successive games are independent.

(a) (i) Calculate an unbiased estimate of p .

(ii) Determine an approximate 95% confidence interval for p . [6]

(b) Brenda also plays this game and she decides to determine a 90% confidence interval for the probability of her winning a game. She therefore plays the game n times and wins x of these games. She correctly calculates an approximate 90% confidence interval to be

[0.455, 0.581]

where the confidence limits are given correct to three decimal places.

Determine

(i) an unbiased estimate of the probability that Brenda wins a game,

(ii) the value of n ,

(iii) the value of x . [7]

4. A motoring organisation wishes to compare the fuel consumption of two car models A and B. It therefore sets up a test in which 50 cars of each model are each supplied with 5 litres of fuel and are driven at a predetermined speed along a track until the fuel is used up. Let x denote the distance (in miles) travelled by each car of model A before stopping and let y denote the distance (in miles) travelled by each car of model B before stopping. The results are summarised below.

$$\sum x = 2565, \quad \sum x^2 = 131659, \quad \sum y = 2590, \quad \sum y^2 = 134232$$

- (a) State suitable hypotheses to investigate whether or not there is a difference between the mean distances travelled by model A and model B cars when given 5 litres of fuel. [1]
- (b) Calculate the p -value of the above results and interpret your value in context. [10]
5. The temperature $y^\circ\text{C}$ in an oven x minutes after switching on the oven can be assumed to satisfy the equation $y = \alpha + \beta x$. In order to estimate α and β , the following measurements were made.

x	0	1	2	3	4	5
y	20.0	34.4	49.3	65.6	79.7	96.5

- (a) Calculate least squares estimates for α and β . [8]
- (b) The values of x are exact but the values of y are subject to independent normally distributed measurement errors with mean zero and standard deviation 0.75. Determine a 99% confidence interval for β . [5]

TURN OVER

6. The probability density function of the continuous random variable X is given by

$$\begin{aligned} f(x) &= \frac{2x}{a^2} && \text{for } 0 \leq x \leq a, \\ f(x) &= 0 && \text{otherwise,} \end{aligned}$$

where a is an unknown positive constant.

(a) Obtain an expression for $E(X)$ and show that

$$\text{Var}(X) = \frac{a^2}{18}. \quad [7]$$

(b) In order to estimate a , a random sample of n observations of X is taken.

(i) The mean of the observations in the sample is denoted by \bar{X} . Find the value of the constant c such that

$$U = c\bar{X}$$

is an unbiased estimator for a and obtain an expression for the variance of U .

(ii) Let Y denote the largest observation in the sample. You are given that

$$E(Y) = \frac{2na}{2n+1} \quad \text{and} \quad \text{Var}(Y) = \frac{na^2}{(n+1)(2n+1)^2}.$$

Find the value of the constant d such that

$$V = dY$$

is an unbiased estimator for a and obtain an expression for the variance of V .

(iii) Show that

$$\frac{\text{Var}(U)}{\text{Var}(V)} = \frac{n+1}{2}.$$

State, with a reason, which is the better estimator, U or V .

[13]