

PH4


| Question | | Marking details | Marks Available |
|-------------------------|-----|---|-----------------|
| 1 | (a) | <p>The total momentum (of a system) is constant (must have total, sum, vector sum etc.) (1)</p> <p>Provided no <u>external</u> [resultant] force (1)</p> <p>e.g. - The momentum before and after a collision is the same - 0</p> <p>When two particles collide the sum of momentum stays the same as long as no forces are involved - 1</p> | 2 |
| | (b) | <p>$\lambda = \frac{c}{f}$ i.e. rearranged or $E = hf = 2.13 \times 10^{-13}$ [J] (1)</p> <p>$p = \frac{h}{\lambda}$ used or $p = \frac{E}{c}$ quoted (1)</p> <p>Final evidence $p = \frac{6.63 \times 10^{-34}}{9.35 \times 10^{-13}}$ or $p = \frac{2.128 \times 10^{-13}}{3 \times 10^8}$ (= 7.09×10^{-22}) (1)</p> | 3 |
| | (c) | <p>Reasonable attempt at cons of mom e.g. initial p of Ni = final p of Ni $\pm p$ of photon (1)</p> <p>$2440 \times 9.95 \times 10^{-26} = 7 \times 10^{-22} - 9.95 \times 10^{-26} v$ (1)</p> <p>Answer = 4700 [m s^{-1}] or slightly different dependent on (b) (1) ecf on p</p> | 3 |
| Question 1 Total | | | [8] |

| Question | | Marking details | Marks Available |
|-------------------------|---------|--|-----------------|
| 2 | (a) | $\frac{1}{3}\rho\overline{c^2}V = nRT$ i.e. some sort of combining both equations (1) Realising $\rho = \frac{Nm}{V}$ (any mistakes in N and m means max of 1/3) (1) (or equivalent steps) Clear algebra with no mistakes leading to $\frac{1}{2}m\overline{c^2} = \frac{3}{2}kT$ or $\frac{1}{2}m\overline{c^2} = \frac{3}{2}\frac{R}{N_A}T$ (if it's difficult to follow don't award the mark - needs to be clear) (1) | 3 |
| | (b) (i) | Mass of argon molecule = $6.3(08)\times 10^{-26}$ (1) Algebra or equivalent method $T = \frac{m\overline{c^2}}{3k}$ or $T = \frac{N_A m\overline{c^2}}{3R}$ (1) Answer = 605 [K] (1) ecf | 3 |
| | (ii) | $\sqrt{\overline{c^2}} \propto \sqrt{T}$ or correct substitution of 1 210 K (ecf) and algebra (1) Answer = $630 \times \sqrt{2}$ or 891 [m s ⁻¹] (1) | 2 |
| Question 2 Total | | | [8] |

| Question | | | Marking details | Marks Available |
|-------------------------|-----|-------|---|-----------------|
| 3 | (a) | (i) | Graph is straight line through origin [hence proportional] (1) (accept acceleration is proportional to displacement) Negative gradient [hence direction ok] (1) | 2 |
| | | (ii) | Gradient calculated correctly i.e. $\frac{1}{0.028}$ or 36 (or k calculated from $ma = kx$ i.e. 7.14 N m^{-1}) (1) Gradient = angular velocity squared i.e. method explained Or $f = \left(2\pi\sqrt{\frac{m}{k}}\right)^{-1}$ i.e. equation for T and $f = 1/T(1)$ Answer = $\frac{5.98}{2\pi} = 0.95 \text{ [Hz]}$ (1) | 3 |
| | | (iii) | 1 m s^{-2} read off graph Or $6^2 \times 0.028 = 1 \text{ [m s}^{-2}\text{]} \text{ etc.}$ | 1 |
| | | (iv) | Max speed = ωA or implied (= 0.167) (1) KE = $\frac{1}{2}mv^2$ or implied (1) Answer = 2.8 [mJ] (1) ecf | 3 |
| | | (v) | $v = A\omega\cos\omega t$ used or $\varepsilon = 0$ stated (1) Rearrangement e.g. $\omega t = \cos^{-1}\frac{v}{A\omega}$ or implied (1) Correct answer = 0.156 [s] (1) ecf | 3 |
| | (b) | | KE to PE or PE to KE (1) PE is both GPE and EPE (1) Energy gradually lost due to friction or air resistance or internal energy of spring/air etc. Not sound, not heat by itself - needs more e.g. lost as heat to the air ok (1) Detail of energy loss e.g. internal energy of air, KE of air particles | 4 |
| Question 3 total | | | | [16] |

| Question | | Marking details | Marks Available |
|-------------------------|-----|--|-----------------|
| 4 | (a) | $T = \frac{pV}{nR}$ seen or implied (1) | 3 |
| | | Evidence of 1 correct substitution (1) | |
| | | Evidence of all 3 substituted correctly (1) A - $(0.500 \pm 0.002, 80\,000 \pm 2\,000)$ B - $(0.260 \pm 0.002, 235\,000 \pm 2\,000)$ C - $(0.260 \pm 0.002, 80\,000 \pm 2\,000)$ | |
| | (b) | $U = \frac{3}{2}nRT$ used (1) | 3 |
| | | Evidence of ΔT being used or differences in U being calculated (once) (1) AB = 31 500 [J], BC = - 60 500 [J], CA = 29 000 [J] (1) | |
| | (c) | (i) AB approximated as a trapezium (accept triangle gives 19 000 J) (1) AB = - 38 000 [J] (1) AB \approx (- 32 000 \pm 3 000) J due to better method ✓✓✓ e.g. two trapezia or 2 triangles or square counting, or any attempt at integrating pV (unlikely) etc. (i.e. 2 marks for good method 1 mark for correct answer) (1) | 3 |
| | | (ii) BC = 0 (independent) | 1 |
| | | (iii) CA = 19 200 [J] | 1 |
| | (d) | (i) $Q = \Delta U + W$ i.e. equation used (1) Correct answer with their figures e.g. $31\,500 - 38\,000 = -6\,500$ (also ecf possible for $31\,500 + 38\,000 = 69\,500$) (1) | 2 |
| | | (ii) No time for heat transfer | 1 |
| Question 4 Total | | | [14] |

| Question | | | Marking details | Marks Available |
|----------|-----|-------------------------|--|-----------------|
| 5 | (a) | (i) | $g = \frac{GM}{r^2}$ used (1) Answer = 3.7 m s^{-2} or N kg^{-1} or equivalent (1) UNIT MARK | 2 |
| | | (ii) | $V_g = -\frac{GM}{r}$ used (1) Answer = $\pm 9.02 \text{ [MJ kg}^{-1}\text{]}$ (1) ecf on km conversion | 2 |
| | | (iii) | Negative amount of work bringing mass from ∞ (accept no work done bringing from ∞ or system will do work or work is done in the other direction etc.) | 1 |
| | (b) | (i) | $PE = V_g \times m$ or implied (1) $KE = \frac{1}{2}mv^2$ used (1) Answer = $656 \text{ kJ} - 4.1 \text{ MJ} = -3.44 \text{ [MJ]}$ (1) | 3 |
| | | (ii) | $-\frac{GMm}{r} = -3.44 \text{ MJ}$ (1) $r = 2905 \text{ km}$ (1) Height = 465 [km] (1) ecf | 3 |
| | | Question 5 Total | | |

| Question | | Marking details | Marks Available |
|----------|-----|---|-----------------|
| 6 | (a) | <p>Arrows pointing towards charges similar to shown ✓✓</p> <p>Arrows pointing away from charges similar to shown ✓</p>  | 2 |
| | (b) | <p>$E = \frac{Q}{4\pi\epsilon_0 r^2}$ used (1)</p> <p>The 2 vertical components cancel or no field into or out of page (1)</p> <p>Pythagoras or trig e.g. $\sqrt{5^2 - 4^2} = 3$ or recognising 3,4,5 triangle (equivalent is to realise $\cos \theta = 3/5$ or $\theta = 53^\circ$ etc.) (1)</p> <p>2 nC charge field x2 and x3/5 ecf (for horizontal components) (1)</p> <p>Calculations all ok e.g. $8\ 640 = 7\ 200 \times 2 \times 3/5$ or equivalent shown (1)</p> | 5 |
| | (c) | <p>$V = \frac{Q}{4\pi\epsilon_0 r}$ used (1)</p> <p>Attempt at adding all 3 potentials (1)</p> <p>- 360 -360 - 432 = - 1 152 V or J C^{-1} or equivalent (1) UNIT MARK</p> | 3 |
| | (d) | <p>Use of $\text{PE} = q\Delta V$ must be a change (1)</p> <p>Rearrangement i.e. $v^2 = \frac{2 \times \text{PE}}{m}$ allow ecf on V (1)</p> <p>Answer = $18.3 \times 10^6 \text{ [m s}^{-1}\text{]}$ (ecf only if a ΔV used) (1)</p> | 3 |
| | | Question 6 Total | [13] |

| Question | | Marking details | Marks Available |
|----------|-----|--|-----------------|
| 7 | (a) | $T = 2\pi \sqrt{\frac{(1.4 \times 10^{10})^3}{6.67 \times 10^{-11} \times (1.6 \times 10^{29} + 3.7 \times 10^{27})}} \quad (1)$ <p>Answer = 3.15×10^6 [s] or implied (3.19×10^6 s if M_2 omitted) (1)</p> <p>36.5 [days] (1) (36.9 if M_2 omitted gets 2/3)</p> | 3 |
| | (b) | (i)&(ii) $r_1 = \frac{M_2}{M_1 + M_2} d \quad \text{used or } M_1 r_1 = M_2 r_2 \quad \text{used} \quad (1)$ <p>Star orbit radius = 0.032×10^{10} [m] (1)</p> <p>Planet orbit radius = 1.37×10^{10} [m] (1)</p> | 3 |
| | (c) | (i) $v = \frac{2\pi r}{T} \quad \text{or} \quad v = \omega r \quad \text{and} \quad \omega = 2\pi f \quad (1)$ $v = \frac{2\pi \times 0.032 \times 10^{10}}{3.15 \times 10^6} (= 631) \quad (1) \quad \text{ecf}$ | 2 |
| | | (ii) $\frac{\Delta\lambda}{\lambda} = \frac{v}{c} \quad \text{values substituted or not possible} \quad (1)$ <p>Answer = 3.9 [pm] because mean radial speed unknown (1)</p> <p>Don't penalise using $2 \times v$ if explained</p> <p>Question 7 Total</p> | 2 |
| | | | [10] |



WJEC
245 Western Avenue
Cardiff CF5 2YX
Tel No 029 2026 5000
Fax 029 2057 5994
E-mail: exams@wjec.co.uk
website: www.wjec.co.uk